

<https://helda.helsinki.fi>

New records of lichens and allied fungi from the Leningrad Region, Russia. IX.

Stepanchikova, Irina S.

2018-08

Stepanchikova , I S , Himelbrant , D , Motiejunaite , J , Ahti , T T , Suija , A , Kuznetsova , E S & Dyomina , A V 2018 , ' New records of lichens and allied fungi from the Leningrad Region, Russia. IX. ' , Folia Cryptogamica Estonica , vol. 55 , pp. 117-124 . <https://doi.org/10.12697/fce.2018.55.12>

<http://hdl.handle.net/10138/273444>

<https://doi.org/10.12697/fce.2018.55.12>

unspecified

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

New records of lichens and allied fungi from the Leningrad Region, Russia. IX

Irina S. Stepanchikova^{1,2}, Dmitry E. Himelbrant^{1,2}, Jurga Motiejūnaitė³, Teuvo Ahti⁴,
Ave Suija⁵, Ekaterina S. Kuznetsova^{1,2}, Aleksandra V. Dyomina¹

¹Department of Botany, St. Petersburg State University, Universitetskaya emb. 7–9, 199034 St. Petersburg, Russia.

E-mails: d_brant@mail.ru, stepa_ir@mail.ru, igel_kuzn@mail.ru, shelka-dyomina@yandex.ru

²Laboratory of Lichenology and Bryology, Komarov Botanical Institute RAS, Professor Popov St. 2,
197376 St. Petersburg, Russia

³Laboratory of Mycology, Institute of Botany, Nature Research Centre, Žaliųjų Ežerų 49, LT–08406 Vilnius, Lithuania.

E-mail: jurga.motiejunaite@botanika.lt

⁴Botanical Museum, Finnish Museum of Natural History, University of Helsinki, P.O. Box 7, FI-00014 Helsinki, Finland.

E-mail: teuvo.ahti@helsinki.fi

⁵Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, Lai St. 40, 51005 Tartu, Estonia.

E-mail: ave.suija@ut.ee

Abstract: Eight species of lichens and seven lichenicolous fungi are reported for the first time for St. Petersburg, the whole Leningrad Region or its western or eastern parts. The lichen *Xylographa septentrionalis* is reported for the first time for Russia, Europe and Asia, the lichenicolous fungus *Skyttea gregaria* is new for Russia, the lichen *Xylographa pallens*, lichenicolous fungi *Didymellopsis collematum* and *Intralichen lichenicola* are new for the North-Western European Russia. The most interesting records are briefly discussed.

Keywords: St. Petersburg, Russia, *Didymellopsis collematum*, *Intralichen lichenicola*, *Skyttea gregaria*, *Xylographa pallens*, *Xylographa septentrionalis*

INTRODUCTION

The present paper continues the series of publications on new and noteworthy findings of lichens and allied fungi from the Leningrad Region and St. Petersburg (see e.g. Kuznetsova et al., 2007, 2012; Stepanchikova et al., 2010, 2011; Himelbrant et al., 2014, 2016, 2017). *Skyttea gregaria* and *Xylographa septentrionalis* are reported for the first time for Russia, *Didymellopsis collematum*, *Intralichen lichenicola* and *Xylographa pallens* are new for the North-Western European Russia. Five species (*Arctomia fascicularis*, *Xanthomendoza weberi*, *Peltigera occidentalis*, *Trichonectria anisospora* and *Usnea florida*) are new for the whole Leningrad Region (including St. Petersburg), three (*Lecidea leprarioides*, *Taeeniolella beschiana* and *Tremella phaeophysciae*) – for St. Petersburg, two (*Dermatocarpon meiohyllizum* and *Lichenodiplis lecanorae*) – for the eastern part of the Leningrad Region.

The revealed lichen diversity of Leningrad Region and St. Petersburg counts now ca. 1090 species, including 940 lichens, 120 lichenicolous and 30 saprobic fungi.

MATERIAL AND METHODS

Specimens were collected by Dmitry E. Himelbrant, Irina S. Stepanchikova, Ekaterina S. Kuznetsova, Aleksandra V. Dyomina and Vladimir A. Plotnikov in 2007–2017 in the eastern and western parts of Leningrad Region or in St. Petersburg, and are deposited in the lichen herbaria of St. Petersburg State University (LECB) and Institute of Botany, Nature Research Centre in Vilnius (BILAS). Furthermore, we investigated several specimens of lichens and lichenicolous fungi kept in the herbarium of University of Helsinki (H). The cited specimens were mainly identified by the authors of the paper, if otherwise, the identifier's name is indicated in the annotation of the species.

The names of the main collectors in the species list are abbreviated as follows: DH – Dmitry E. Himelbrant, EK – Ekaterina S. Kuznetsova, IS – Irina S. Stepanchikova, AD – Aleksandra V. Dyomina. The subdivision of the Leningrad Region (LR) was published in our previous paper (Stepanchikova et al., 2010); the following abbreviations have been used here: ELR – Eastern

Leningrad Region, SPb – St. Petersburg, WLR – Western Leningrad Region. The biogeographical provinces of Eastern Fennoscandia are abbreviated traditionally (Kotiranta et al., 1998): Ik – Isthmus karelicus, Ka – Karelia australis, Kol – Karelia olonetsensis. All geographical coordinates are given in the spatial reference system WGS 1984. Lichenicolous fungi are marked with #. The nomenclature of taxa generally follows Nordin et al. (2011) and Randlane et al. (2016); for the species not mentioned in the checklists, special papers are cited (Fedorenko et al., 2012; Spribille et al., 2014).

THE SPECIES

ARCTOMIA FASCICULARIS (L.) Otálora & Wedin – WLR, Kingisepp District, Kotelsky protected area, NE shore of Lake Babinskoe, SE of Savikino, 59°36'07.1"N, 28°36'01.9"E, alt. 20 m, old-growth birch-aspen-black alder forest, on bark of *Populus tremula* L., 15.04.2007, leg. DH (H s. n.); ELR, Podporozhje District, NE of Rodionovo, bank of the Kuzra River, 60°57'16"N, 35°10'34"E, old-growth spruce forest (age of spruces 180–200 years) with *Sphagnum* mosses, old pines, sparse aspens and birches, on bark of *P. tremula*, 03.10.2014, leg. DH & IS (LECB); ELR, Boksitogorsk District, W of Somino, between lakes Loshevo and Lachta, 59°20'42"N, 34°49'59"E, alt. 167 m; spruce forest with old aspens and herbs on the slope near the road, on bark of *P. tremula*, 18.07.2011, leg. IS (H s. n.). – New to LR. Distribution in North-Western European Russia outside of LR: Republic of Karelia (Tarasova & Stepanchikova, 2016). Distribution in Fennoscandia and Baltic countries: Norway, Sweden (Nordin et al., 2011). Characterized by subfoliose thallus forming thick (and swelling when wet) cushions to 2–3 cm diam., membranaceous and fenestrate, with numerous accessory lobules (Jørgensen, 2007). Apothecia should be common, but not in material from LR.

BACIDINA SULPHURELLA (Samp.) M. Hauck & V. Wirth – SPb, Petrodvorets District, Strel'na, forest between the Strelka River and Novye Zavody Street, 59°50'02.9"N, 29°59'11.4"E, pine forest with dense rowan undergrowth and young oaks, on bark of *Sorbus aucuparia* L., 03.06.2017, leg. DH & IS (LECB). – Previously reported from SPb (Stepanchikova et al., 2015), but the identification was revised later (Himelbrant et al., 2017).

Known from WLR (Stepanchikova et al., 2017). Distribution in North-Western European Russia outside LR and SPb: not recorded. Distribution in Fennoscandia and Baltic countries: Estonia (Randlane et al., 2016), Lithuania (Motiejūnaitė, 2017).

DERMATOCARPON MEIOPHYLLIZUM Vain. – ELR, Kol, Podporozhje District, pier on the Vazhinka River in the village Grishino, 61°05'11.6"N, 34°05'12.8"E, on siliceous boulders in the river, 07.07.2016, leg. IS & AD (LECB). – New to ELR, known from SPb (Himelbrant et al., 2016). Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007). Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), probably Estonia (Randlane et al., 2016: doubtful record).

DIDYMELLOPSIS COLLEMATUM (J. Steiner) Grube & Hafellner – ELR, Ka, Vyborg District, town of Vyborg (former Viipuri), [60°43'N, 28°45'E], on thallus of *Enchylium bachmanianum* (Fink) Otálora et al. growing on terricolous mosses, 1851, leg. W. Nylander (H 6022460). – New to North-Western European Russia, in Russia known from Krasnoyarsk Territory (Zhurbenko & Santesson, 1996). Distribution in Fennoscandia and Baltic countries: Estonia (Randlane et al., 2016). Characterized by perithecioid ascocarps (pseudothecia) 0.2–0.3 mm diam., brown-pigmented dark exciple, 8-spored fissitunicate asci 80–100 × 13–15 µm, 1-septate hyaline ascospores 20–26 × 5–10 µm and inhabiting *Collema* s. lat. Another *Didymellopsis* known to occur on *Collema* s. lat. and *Leptogium* s. lat., *D. pulposi* (Zopf) Grube & Hafellner, is characterized by smaller ascomata (0.17–0.2 mm diam.), shorter ascospores (13–22 × 5–7 µm), and 4–8-spored, shorter asci (40–90 × 10.5–13.5 µm) (Grube & Hafellner, 1990). Our specimen had 8-spored asci and ascospores measuring 20–23 × 5.5–7 µm.

FELLHANERA GYROPHORICA Sérus., Coppins, Diedrich & Scheid. – WLR, Kingisepp District, 0.2 km S of Lake Kopanskoe, S to the highway, 59°42'02.8"N, 28°42'10.8"E, old-growth black alder-birch-spruce forest, on bark of old *Picea* sp., 13.04.2007, leg. DH (H s. n.). – We previously reported this specimen from the Leningrad Region as the only from Russia (Handbook..., 2008), but without collecting data (exact locality etc.). Distribution in Fennoscandia and Baltic countries: Estonia (Randlane et al., 2016).

Characterized by greenish matt thallus consisting of goniocysts, round or irregular-shaped rose-brown apothecia 0.3–0.7 mm diam. (absent in our material); pycnidia numerous, very distinctive – 0.1–0.25 mm diam., sessile to shortly stipitate, rose to light-brown, with widely open ostiole and simple pycnospores $(2.5)3\text{--}3.5(4.5) \times (0.7)1\text{--}1.5 \mu\text{m}$, contain gyrophoric acid (Handbook..., 2008).

INTRALICHEN LICHENICOLA (M.S. Christ. & D. Hawksw.) D. Hawksw. & M. S. Cole – SPb, Petrodvorets District, Strel'na, proposed protected area 'Strelka River Valley', between the Strelka River and Novye Zavody Street, $59^{\circ}50'13.6''\text{N}$, $30^{\circ}00'04.7''\text{E}$, anthropogenic wasteland, on apothecia of *Candelariella* cf. *vitellina* (Hoffm.) Müll. Arg. on concrete post, 04.06.2017, leg. DH & IS (BILAS 10928). – New to North-Western European Russia, the nearest localities in Russia belong to Tver' Region (Notov et al., 2011). Distribution in Fennoscandia and Baltic countries: Norway, Sweden (Nordin et al., 2011), Lithuania (Motiejūnaitė, 2017). Dematiaceous hyphomycetous fungus with immersed mycelium dispersed inside the host hymenium and turning apothecial disc black (Fig. 1) and forming conidiophores on its surface; conidia pale to dark brown, very variable in shape, multicellular ($18\text{--}25 \times 6\text{--}12 \mu\text{m}$), arising as elongated chains of cells (Hawksworth, 1979). Our specimen was in concurrence with the description in the protologue.




Fig. 1. Apothecia of *Candelariella* cf. *vitellina* darkened by infection of *Intralichen lichenicola* (BILAS 10928). Scale bar = 1 mm.

LECIDEA LEPRARIOIDES Tønsberg – SPb, Ik, Kurortny District, Komarovo (former Kellomäki), nature monument 'Komarovskiy Bereg', W of the railway station Komarovo, $60^{\circ}11'17''\text{N}$, $29^{\circ}46'02''\text{E}$, pine forest with *Calluna vulgaris* L., on bark of *Pinus sylvestris* L., 07.05.2017, leg. DH & IS (LECB). – New to SPb, known from ELR (Kuznetsova et al., 2007) and WLR (Stepanchikova et al., 2013b). Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007), Novgorod Region (Stepanchikova et al., 2013a). Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), Estonia (Randlane et al., 2016).

LICHENODIPLIS LECANORAE (Vouaux) Dyko & D. Hawksw. – ELR, Kol, Podporozhje District, proposed protected area 'Severo-Svirsky', NW of Soginitsy, right side of the Svyatukha River, $61^{\circ}13'06.6''\text{N}$, $33^{\circ}56'13.6''\text{E}$, old-growth spruce forest with *Sphagnum* spp., on apothecia and thallus of *Mycoblastus alpinus* (Fr.) Th. Fr. ex Hellb. on bark of *Picea* sp. (branches), 03.07.2016, leg. AD & IS (BILAS). – New to ELR, known from WLR (Stepanchikova et al., 2011). Distribution in North-Western European Russia outside of LR: not reported. Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), Estonia (Randlane et al., 2016), Latvia (Motiejūnaitė et al., 2016), Lithuania (Motiejūnaitė, 2017).

PELTIGERA OCCIDENTALIS (E. Dahl) Kristinsson – WLR, Priozersk District, S to Lake Parijärvi (Dvinskoe), ca 6 km S of Hiitola, $[61^{\circ}09'\text{N}$, $29^{\circ}47'\text{E}]$, on soil, 01.08.1935, leg. M. Laurila, det. O. Vitikainen (H s. n.). – New to LR. Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007). Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011). The species resembles *P. neopolydactyla* (Gyeln.) Gyeln., from which it differs by having thicker, rigid thallus which is greenish when moist (Vitikainen, 2007).

PELTIGERA VENOSA (L.) Hoffm. – WLR, Luga District, proposed protected area 'Jaschera-Lemovzha', valley of the Jaschera River, $58^{\circ}53'45.9''\text{N}$, $29^{\circ}50'33.3''\text{E}$, steep slope of deep river valley, on sandstone, 11.05.2017, leg. IS & Vladimir A. Plotnikov (LECB). – First record in WLR after 1935 (Räsänen, 1939), and probably the only locality known now in LR. In 1994 the species was collected in ELR by Orvo Vitikainen

(Kuznetsova et al., 2007), but we did not find it at the same locality (Cape Podschel'je) in course of our later field studies. Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007), Pskov Region (Istomina & Likhacheva, 2010). Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), Estonia (Randlane et al., 2016), Latvia (Āboliņa et al., 2015), Lithuania (Motiejūnaitė, 2017: extinct).

SKYTTEA GREGARIA Sherwood, D. Hawksw. & Coppins – WLR, Ik, Priozersk District, E part of Konevets Island, SW of Cape Rodushka, 60°51'00.5"N, 30°36'58.9"E, old-growth spruce forest with *Vaccinium myrtillus* L. and patches of *Sphagnum* spp., on thallus of *Violella fucata* (Stirt.) T. Sprib. growing on twig of *Picea abies*, 04.08.2017, leg. EK, IS & DH (BILAS 10921); ELR, Kol, Podporozhje District, proposed protected area 'Severo-Svirsky', NW of Soginitsy, right side of the Svyatukha River, 61°13'13.4"N, 33°55'56.3"E, old-growth spruce forest with *Sphagnum* spp., on thallus of *V. fucata* on bark of *Picea* sp., 04.07.2016, leg. AD & IS (BILAS); 80 m W to Lake Maloe Kujvozero and 100 m W to the railway, 61°07'01.8"N, 34°23'44.3"E, in wet old-growth spruce forest with *Sphagnum* spp., on thallus of *V. fucata* on twigs of *Picea* sp., 02.07.2016, leg. IS & AD (BILAS 10923). New to Russia. Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), Estonia (Randlane et al., 2016). Apparently a host-specific lichenicolous fungus, characterized by dark-greyish apothecia, at first immersed, becoming erumpent, 0.1–0.2 mm in diameter, with urceolate disc, striate inrolled margin when dry (Fig. 2), with greenish exciple which does not react K+ violet and which is covered with olive-brown hairs (visible under microscope), 5–20 µm long. Asci 1–, 8-spored; spores ovoid, simple, 7–8.5 × 3–3.5(–4) µm (Sherwood et al., 1980; Diederich & Etayo, 2000).

TAENIOLELLA BESCHIANA Diederich – SPb, Ik, Kurortny District, Komarovo (Kellomäki), nature monument 'Komarovskiy Bereg', W of the railway station Komarovo, 60°11'17"N, 29°46'02"E, pine forest with *Calluna vulgaris*, on thallus of *Cladonia* sp. (Fig. 3) on soil, 07.05.2017, leg. DH & IS (BILAS 10927). – New to SPb, known from WLR (Himmelbrant et al., 2017). Distribution in North-Western European Russia outside of LR:




Fig. 2. Ascomata of *Skyttea gregaria* on thallus of *Violella fucata* (BILAS 10921). Scale bar = 0.2 mm.




Fig. 3. Mycelium of *Taeniolella beschiana* on thallus of *Cladonia* sp. (BILAS 10927). Scale bar = 0.5 mm.

Republic of Karelia (Fadeeva et al., 2007). Distribution in Fennoscandia and Baltic countries: Norway, Sweden (Nordin et al., 2011), Estonia (Randlane et al., 2016), Lithuania (Motiejūnaitė, 2017).

TREMELLA PHAEOPHYSCIAE Diederich & M. S. Christ. – SPb, Pushkin District, nature monument 'Outcrops on the Popovka River', Popovka River valley, SW of Pavlovsk railway station, 59°39'37"N, 30°22'09"E, grey alder-aspens floodplain forest with shrubs, on thallus of *Phaeophyscia orbicularis* (Neck.) Moberg on bark of *Padus*

avium Mill., 08.05.2010, leg. DH & EK (LECB); SPb, Petrodvorets District, Strel'na, proposed protected area 'Strelka River Valley', left bank of the Strelka River, 59°50'09.0"N, 30°01'16.1"E, old willows with *Padus avium* on the meadow, on thallus of *Phaeophyscia orbicularis* on bark of *Padus avium*, 04.06.2017, leg. DH & IS (LECB). – New to SPb, known from WLR (Himmelbrant et al., 2016). Distribution in North-Western European Russia outside of LR: not recorded. Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011), Estonia (Randlane et al., 2016), Latvia (Motiejūnaitė et al., 2016), Lithuania (Motiejūnaitė, 2017).

TRICHONECTRIA ANISOSPORA (Lowen) van den Boom & Diederich – ELR, Kol, Podporozhje District, proposed protected area 'Severo-Svirsky', N of Lake Vachozero, 61°04'16.2"N, 34°20'21.2"E, birch forest with young spruces, on thallus of *Hypogymnia physodes* (L.) Nyl. on spruce branches, 06.20.2017, leg. DH & IS (LECB); same place, 61°03'30.8"N, 34°20'50.8"E, aspen-spruce forest, on thallus of *H. physodes* on spruce branches, 06.20.2017, leg. DH & IS (LECB). – New to LR. Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007). Distribution in Fennoscandia and Baltic countries: Norway, Sweden (Nordin et al., 2011), Finland (Stenroos et al. 2016), Estonia (Randlane et al., 2016), Latvia (Motiejūnaitė et al., 2016), Lithuania (Motiejūnaitė, 2017). Characterized by orange setose perithecia, initially immersed, later superficial, 1-septate spores, strongly constricted at septum, with unequal cells, and by variable ascospore form and size, even in the same ascus (Lowen, 1989). The fungus is narrowly host-specific, causes characteristic blue-green, clearly bordered infection patches on upper surface of host thallus and apparently is spreading lately (Brackel, 2006; Motiejūnaitė et al., 2011).

USNEA FLORIDA (L.) F. H. Wigg. – WLR, Luga District, proposed protected area 'Jaschera-Lemovzha', valley of the Jaschera River, 58°53'46.4–46.9"N, 29°50'36.2–45.2"E, elm-linden-maple-spruce forest in deep river valley, on bark of *Ulmus* sp., *Acer platanoides* L. and *Tilia cordata* Mill. (branches), 11.05.2017, leg. Vladimir A. Plotnikov & IS (H s. n., LECB). – First reliable record from LR. The species was reported from LR and SPb several times in the past, latest in 1922 (Elenkin, 1922), but no correctly identified

specimens are known. Distribution in North-Western European Russia outside of LR: Pskov Region (Istomina & Likhacheva, 2010). Distribution in Fennoscandia and Baltic countries: Norway, Sweden (Nordin et al., 2011), Latvia (Āboliņa et al., 2015), Lithuania (Motiejūnaitė, 2017: extinct). Morphologically very distinctive abundantly fertile shrubby species with large fibrillose apical apothecia containing alecortic acid, and with papillose and fibrillose main branches while soralia or isidiomorphs are absent (Smith et al., 2009; Clerc, 2011).

XANTHOMENDOZA WEBERI (S.Y. Kondr. & Kärnefelt) L. Lindblom – WLR, Ka, Vyborg District, ca 1.5 km NEE of Sovetsky (former Rokkala near St. Johannes), [60°33'N, 28°43'E], on bark of *Betula* sp., 20.04.1895, leg. Bertil R. Poppius, det. Sergii Y. Kondratyuk, 2007 (H 8003826). – New to LR. Distribution in North-Western European Russia outside of LR: Republic of Karelia (Fadeeva et al., 2007). Distribution in Fennoscandia and Baltic countries: Latvia (Motiejūnaitė et al., 2016). In the discussed region could be mistaken with *X. fulva* (Hoffm.) Søchting, Kärnefelt & S.Y. Kondr., from which differs by longer, more branched lobes and presence of helmet-shaped soralia (Handbook..., 2004). The material collected by Poppius partly has helmet-shaped soralia and is in good accordance with the description (Kondratyuk & Kärnefelt, 2003).

XYLOGRAPHA PALLENS (Nyl.) Malmgren – ELR, Ik, Priozersk District, Torfyanoe (former Konnitsa, Pyhäjärvi), NW vicinity of Lake Komsomol'skoe (Kiimajärvi), [60°51'N, 29°56'E], on lignum, 07.1897, leg. Carl Gustaf Wilhelm Lång, det. Toby Spribille [H 8003830, previously identified as *X. parallela* (Ach.: Fr.) Fr.]. – New to North-Western European Russia, previously reported from Murmansk Region (Spribille et al., 2014). Distribution in Fennoscandia and Baltic countries: Norway, Sweden, Finland (Nordin et al., 2011; Spribille et al., 2014). Characterized by ascomata with lateral growth of parallela-type up to 2.8 mm long, clustered, forming 'stars'; thallus immersed to superficial, containing stictic acid as major substance (Spribille et al., 2014).

XYLOGRAPHA SEPTENTRIONALIS T. Sprib. – ELR, Ik, Kirillovskoe (Muolaa, Perkjärvi), [60°28'N, 29°20'E], on wood, 23.10.1893, leg. Alfred Oswald Kihlman (Kairamo), det. Toby Spribille [H 8004005, previously identified as *X. vitiligo*

(Ach.) J.R. Laundon]. – New to Russia and Europe, described from Canada (Spribille et al., 2014). We also report the species here for the first time for Asia: several specimens from basin of the Yenisei River by M. Brenner are deposited in H [det. Toby Spribille, H s. n., previously identified as *X. spilomatica* (Anzi) Th. Fr. = *X. vitiligo* and *X. abietina* (Pers.) Zahlbr. = *X. parallela*]: Krasnoyarsk Territory ('Guv. Jenisejsk'), former settlement Tolsty Nos [70°10'N, 83°12'E], on wood, 31.08.1876, 27.08.1876 and 04.09.1876, leg. M. Brenner (1560 h-A, 18e and 685j, respectively); Dudinka [69°25'N, 86°10'E], on wood, 07.08.1876, leg. M. Brenner (1657g); 'Polovinka' (basin of the Polovinka River), on wood, 15.09.1876, leg. M. Brenner (504d); Nikandrovskie islands [70°40'N, 83°14'E], on wood, 17.08.1876, leg. M. Brenner (1838b1). Similar to *X. vitiligo*, but the soralia and gonioscysts are very small and often blackish, so that it is easily overlooked (Fig. 4); contains stictic and norstictic or only norstictic acid. Apothecia, when present, resemble those of *X. parallela* (Spribille et al., 2014).




Fig. 4. Thallus and apothecia of *Xylographa septentrionalis* (H 8004005). Scale bar = 1 mm. Photo by Saara Velmala.

Taxa excluded from the list of LR:

Lecidea plana (J. Lahm) Nyl. – reported by Vainio (1934); the specimens (H 8005263, 8005264) belong to *Lecidea lapicida* (Ach.) Ach.

ACKNOWLEDGEMENTS

Authors would like to thank Toby Spribille (University of Alberta, Canada) for identification of *Xylographa* spp., and Orvo Vitikainen (University of Helsinki, Finland) for identification of *Peltigera occidentalis*. We are also grateful to

secondary school student Vladimir A. Plotnikov (Academic Gymnasium of St. Petersburg State University, Russia), who participated in field investigations in the Jaschera River valley, Saara Velmala (University of Helsinki, Finland) and Jaan Liira (University of Tartu, Estonia) for assistance in microphotographing. We express our gratitude to anonymous reviewer and Tiina Randlane (University of Tartu) for valuable corrections. The study was carried out within the framework of the institutional research project "The lichen flora of the Russian Federation" (no. AAAA-A18-118031590042-0) of the Komarov Botanical Institute of the Russian Academy of Sciences. Specimens from Krasoyarsk Territory were revised in frame of the project no. 18-05-60093 of Russian Fund for Basic Research (RFBR).

REFERENCES

- Āboliņa, A., Piterāns, A. & Bамbe, B. 2015. *Lichens and bryophytes in Latvia. Checklist*. Salaspils, Latvijas Valsts mežzinātnes institūts "Silava", DU AA Saule. 213 pp. (In Latvian, English introductions).
- Brackel, W. v. 2006. Zur Verbreitung von *Trichonectria anisospora* (Lowen) P. Boom & Diederich. *Meylania* 37: 5–7.
- Clerc, P. 2011. *Usnea*. – In: Thell, A. & Moberg, R. (eds). *Nordic lichen flora 4*. Uppsala University & Nordic Lichen Society, Uppsala. Pp. 107–127.
- Diederich, P. & Etayo, J. 2000. A synopsis of the genera *Skyttea*, *Llimoniella* and *Rhymbocarpus* (lichenicolous Ascomycota, Leotiales). *Lichenologist* 32(5): 423–485. <https://doi.org/10.1006/lich.2000.0290>
- Elenkin, A. A. 1922. *Lichens as an object of pedagogy and scientific research*. St. Petersburg. 132 pp. (In Russian).
- Fadeeva, M. A., Golubkova, N. S., Vitikainen, O. & Ahti, T. 2007. *Conspectus of lichens and lichenicolous fungi of the Republic of Karelia*. Petrozavodsk. 194 pp. (In Russian, English summary).
- Fedorenko, N. M., Stenroos, S., Thell, A., Kärnefelt, I., Elix, J. A., Hur, J.-S. & Kondratyuk, S.Y. 2012. Molecular phylogeny of xanthorioid lichens (Telo-schistaceae, Ascomycota), with notes on their morphology. *Bibliotheca Lichenologica* 108: 45–64.
- Grube, M. & Hafellner, J. 1990. Studien an flechtenbewohnenden Pilzen der Sammelgattung *Didymella* (Ascomycetes, Dothideales). *Nova Hedwigia* 51: 283–360.
- Handbook of the lichens of Russia*. Vol. 10. 2008. St. Petersburg. 515 pp. (In Russian).
- Hawksworth, D. L. 1979. The lichenicolous Hyphomycetes. *Bulletin of the British Museum (Natural History), Botany Series* 6: 183–300.

- Himelbrant, D. E., Motiejūnaitė, J., Stepanchikova, I. S. & Tagirdzhanova, G. M. 2014. New records of lichens and allied fungi from the Leningrad Region, Russia. V. *Folia Cryptogamica Estonica* 51: 49–55. <https://doi.org/10.12697/fce.2014.51.04>
- Himelbrant, D. E., Stepanchikova, I. S., Motiejūnaitė, J., Gagarina, L. V. & Dyomina, A. V. 2016. New records of lichens and allied fungi from the Leningrad Region, Russia. VII. *Folia Cryptogamica Estonica* 53: 25–34. <https://doi.org/10.12697/fce.2016.53.04>
- Himelbrant, D. E., Stepanchikova, I. S., Motiejūnaitė, J., Gerasimova, J. V., Kuznetsova, E. S., Dyomina, A. V. & Tsurykau, A. G. 2017. New records of lichens and allied fungi from the Leningrad Region, Russia. VIII. *Folia Cryptogamica Estonica* 54: 63–70. <https://doi.org/10.12697/fce.2017.54.11>
- Istomina, N. B. & Likhacheva, O. V. 2010. The preliminary list of lichens of the Pskov Region (Russia). (In Russian, English summary). *Novitates Systematicae Plantarum Non Vascularium* 44: 171–199.
- Jørgensen, P. M. 2007. Collemataceae. – In: *Nordic Lichen Flora. Volume 3. Cyanolichens*. Nordic Lichen Society, Uddevalla. Pp. 14–42.
- Kondratyuk, S. & Kärnefelt, I. 2003. Five new Xanthorias from Holarctic. *Ukrainian Botanical Journal* 60: 121–130.
- Kotiranta, H., Uotila, P., Sulkava, S. & Peltonen, S.-L. (eds). 1998. *Red Data Book of East Fennoscandia*. Helsinki. 351 pp.
- Kuznetsova, E., Ahti, T. & Himelbrant, D. 2007. Lichens and allied fungi of the Eastern Leningrad Region. *Norrinia* 16: 1–62.
- Kuznetsova, E. S., Motiejūnaitė, J., Stepanchikova, I. S., Himelbrant, D. E. & Czarnota, P. 2012. New records of lichens and allied fungi from the Leningrad Region, Russia. III. *Folia Cryptogamica Estonica* 49: 31–37.
- Lowen, R. 1989. Two new species of *Nectriella* and an *Acremonium* anamorph. *Memoirs of the New York Botanical Garden* 49: 243–252.
- Motiejūnaitė, J., 2017. Supplemented checklist of lichens and allied fungi of Lithuania. *Botanica Lithuanica* 23(2): 89–106. <https://doi.org/10.1515/botlit-2017-0011>
- Motiejūnaitė, J., Brackel, W. v., Stončius, D., Preikša, Ž., 2011. Contribution to the Lithuanian flora of lichens and allied fungi. III. *Botanica Lithuanica* 17(1): 39–47.
- Motiejūnaitė, J., Chesnokov, S. V., Czarnota, P., Gagarina, L. V., Frolov, I., Himelbrant, D., Konoreva, L. A., Kubiak, D., Kukwa, M., Moisejevs, R., Stepanchikova, I., Suija, A., Tagirdzhanova, G., Thell, A. & Tsurykau, A. 2016. Ninety-one species of lichens and allied fungi new for Latvia with a list of additional records from Kurzeme. *Herzogia* 29(1): 143–163. <https://doi.org/10.13158/heia.29.1.2016.143>
- Nordin, A., Moberg, R., Tønsgberg, T., Vitikainen, O., Dalsätt, Å., Myrdal, M., Snitting, D. & Ekman, S. 2011. *Santesson's Checklist of Fennoscandian Lichen-forming and Lichenicolous Fungi*. Ver. April 29, 2011 – <http://130.238.83.220/santesson/home.php> (15 March 2018).
- Notov, A. A., Himelbrant, D. E. & Urbanavichus, G. P. 2011. *The list of lichens and allied fungi of Tver Region*. (In Russian). Tver. 124 pp.
- Randlane, T., Saag, A. & Suija, A. 2016. *Lichenized, lichenicolous and allied fungi of Estonia*. Ver. December 31, 2016 – <http://esamba.bo.bg.ut.ee/checklist/est/home.php> (15 March 2018).
- Räsänen, V. 1939. Die Flechtenflora der nördlichen Küstengegend am Laatokka-See. *Annales Botanici Societatis Zoologicae-Botanicae Fennicae «Vanamo»* 12(1): 1–240.
- Sherwood, M. A., Hawksworth, D. L. & Coppins, B. J. 1980. *Skyttea*, a new genus of odontotremoid lichenicolous fungi. *Transactions of the British Mycological Society* 75: 479–490. [https://doi.org/10.1016/S0007-1536\(80\)80130-6](https://doi.org/10.1016/S0007-1536(80)80130-6)
- Smith, C. W., Aptroot, A., Coppins, B. J., Fletcher, A., Gilbert, O. L., James, P. W. & Wolseley, P. A. (eds). 2009. *The lichens of Great Britain and Ireland*. The British Lichen Society, London. 1046 pp.
- Spribile, T., Resl, P., Ahti, T., Pérez-Ortega, S., Mayrhofer, H. & Lumbsch, H. T. 2014. Molecular systematics of the wood-inhabiting, lichen-forming genus *Xylographa* (Baeomycetales, Ostropomycetidae) with eight new species. *Symbolae Botanicae Upsalienses* 37 (1): 1–87.
- Stenroos, S., Velmala, S., Pykälä, J. & Ahti, T. 2016. Lichens of Finland. *Norrinia* 30: 1–896.
- Stepanchikova, I. S., Andreev, M. P., Himelbrant, D. E., Motiejūnaitė, J., Schiefelbein, U., Konoreva, L. A. & Ahti, T. 2017. The lichens of Bolshoy Tuters Island (Tytärsaari), Leningrad Region, Russia. *Folia Cryptogamica Estonica* 54: 95–116. <https://doi.org/10.12697/fce.2017.54.14>
- Stepanchikova, I. S., Gagarina, L. V. & Kataeva, O. A. 2013a. New records of lichens and allied fungi from the Novgorod Region, Russia. *Folia Cryptogamica Estonica* 50: 49–55. <https://doi.org/10.12697/fce.2013.50.07>
- Stepanchikova, I. S., Himelbrant, D. E., Dyomina, A. V. & Tagirdzhanova, G. M. 2015. The lichens and allied fungi of the Zapadny Kotlin protected area and its vicinities (Saint Petersburg). *Novitates Systematicae Plantarum Non Vascularium* 49: 265–281.
- Stepanchikova, I. S., Kukwa, M., Kuznetsova, E. S., Motiejūnaitė, J. & Himelbrant, D. E. 2010. New records of lichens and allied fungi from the Leningrad Region. *Folia Cryptogamica Estonica* 47: 77–84.
- Stepanchikova, I. S., Tagirdzhanova, G. M. & Himelbrant, D. E. 2013b. The lichens and allied fungi of the Smorodinka River valley (Leningrad Region). *Novitates Systematicae Plantarum Non Vascularium* 47: 262–278.

- Stepanchikova, I. S., Schiefelbein, U., Alexeeva, N. M., Ahti, T., Kukwa, M., Himmelbrant, D. E. & Pykälä, J. 2011. Additions to the lichen biota of Berezovye Islands, Leningrad Region, Russia. *Folia Cryptogamica Estonica* 48: 95–106.
- Tarasova, V. N. & Stepanchikova I. S. 2016. New lichens to Karelia Republic. *Proceedings of Petrozavodsk State University. Biological sciences* 4 (157): 78–82. (In Russian).
- Vainio, E. A. Lichenographia Fennica IV. Lecideales II. 1934. *Acta Societatis pro Fauna et Flora Fennica* 57(2): 1–592.
- Vitikainen, O. 2007. Peltigeraceae. – In: *Nordic Lichen Flora. Volume 3. Cyanolichens*. Nordic Lichen Society, Uddevalla. Pp. 113–131.
- Zhurbenko, M. P. & Santesson, R. 1996. Lichenicolous fungi from the Russian Arctic. *Herzogia* 12: 147–161.